

FIG.1

- ▲ 0.3 mg/kg, INTRAMUSCULAR ADMINISTRATION, ADMINISTRATION AREA
- 1 mg/kg, INTRAMUSCULAR ADMINISTRATION, ADMINISTRATION AREA
- 3 mg/kg, INTRAMUSCULAR ADMINISTRATION, ADMINISTRATION AREA
- △ 0.3 mg/kg, INTRAMUSCULAR ADMINISTRATION, CONTRALATERAL MUSCLE
- 1 mg/kg, INTRAMUSCULAR ADMINISTRATION, CONTRALATERAL MUSCLE
- 3 mg/kg, INTRAMUSCULAR ADMINISTRATION, CONTRALATERAL MUSCLE
- * 600 µg/kg/hr, CONTINUOUS INTRAVENOUS ADMINISTRATION, 3 HR




FIG.2

- ▲ 0.3 mg/kg, INTRAMUSCULAR ADMINISTRATION
- 1 mg/kg, INTRAMUSCULAR ADMINISTRATION
- 3 mg/kg, INTRAMUSCULAR ADMINISTRATION
- △ 6 µg/kg/hr, CONTINUOUS INTRAVENOUS ADMINISTRATION, 3 HR
- 60 µg/kg/hr, CONTINUOUS INTRAVENOUS ADMINISTRATION, 3 HR
- 600 µg/kg/hr, CONTINUOUS INTRAVENOUS ADMINISTRATION, 3 HR
- * 0.3 mg/kg, INTRAVENOUS BOLUS ADMINISTRATION

BEST AVAILABLE COPY




FIG.3

- ▲— 0.3 mg/kg, INTRAMUSCULAR ADMINISTRATION
- 1 mg/kg, INTRAMUSCULAR ADMINISTRATION
- 3 mg/kg, INTRAMUSCULAR ADMINISTRATION
- △— 6 µg/kg/hr, CONTINUOUS INTRAVENOUS ADMINISTRATION, 3 HR
- 60 µg/kg/hr, CONTINUOUS INTRAVENOUS ADMINISTRATION 3 HR
- 600 µg/kg/hr, CONTINUOUS INTRAVENOUS ADMINISTRATION 3 HR
- ×— 0.3 mg/kg, INTRAVENOUS BOLUS ADMINISTRATION




FIG.4

- ▲— 0.3 mg/kg, INTRAMUSCULAR ADMINISTRATION
- 1 mg/kg, INTRAMUSCULAR ADMINISTRATION
- 3 mg/kg, INTRAMUSCULAR ADMINISTRATION
- △— 6 µg/kg/hr, CONTINUOUS INTRAVENOUS ADMINISTRATION, 3 HR
- 60 µg/kg/hr, CONTINUOUS INTRAVENOUS ADMINISTRATION 3 HR
- 600 µg/kg/hr, CONTINUOUS INTRAVENOUS ADMINISTRATION 3 HR
- ×— 0.3 mg/kg, INTRAVENOUS BOLUS ADMINISTRATION




FIG.5





FIG.6

REPLACEMENT SHEET

4/7



BEST AVAILABLE COPY

1: NON-TREATED,
 2: 5 MIN AFTER ADMINISTRATION OF 0.03 mg/kg HGF,
 3: 30 MIN AFTER ADMINISTRATION OF 0.03 mg/kg HGF,
 4: 5 MIN AFTER ADMINISTRATION OF 0.3 mg/kg HGF,
 5: 30 MIN AFTER ADMINISTRATION OF 0.3 mg/kg HGF,
 6: 5 MIN AFTER ADMINISTRATION OF 3 mg/kg HGF,
 7: 30 MIN AFTER ADMINISTRATION OF 3 mg/kg HGF,
 PY: BLOTTING BY ANTI-PHOSPHOTYROSINE ANTIBODY
 c-Met: BLOTTING BY ANTI-c-Met ANTIBODY
 ARROWS SHOW LOCATIONS OF c-Met.

FIG.7




FIG.8




FIG.9




FIG.10




FIG.11

09, 762,188

BEST AVAILABLE COPY




FIG.12




FIG.13